

## Data and References

This document describes how data are used for the guidelines revision process, and lists selected references for each working group. As described in the document *ASCCP Risk-Based Management Consensus Guidelines: New Recommendations for Public Comment*, we use the strongest data available for each management question. Large prospective, longitudinal databases are considered the highest level of evidence, and are used whenever possible. Literature review is used to address questions that cannot be answered by primary data. For areas where neither primary data nor literature provide sufficient evidence, previous guidelines or consensus opinion are used.

**Large prospective, longitudinal databases:** KPNC contains longitudinal data on >1 million women screened triennially from 2003 with Pap/HPV cotesting. Primary risk estimates are calculated using this data set. To confirm that thresholds are appropriate in the larger US setting, confirmation of risk estimates will be performed in several additional datasets, including those from other geographic areas of the US as well as those targeting low-income, minority, and underserved women.

**Literature review:** For questions where primary data are insufficient, systematic literature searches are performed using MESH search terms relative to the disease, management, and/or test concepts. Literature searches focus on material published after the 2012 guidelines.

## Selected references from each working group

*Please note these are not complete lists of all the references used, but curated lists of selected references.*

## Treatment

Arbyn M, Redman CW, Verdoodt F, Kyrgiou M, Tzafetas M, Ghaem-Maghani S, Petry KU, Leeson S, Bergeron C, Nieminen P, Gondry J, Reich O, Moss E. Incomplete excision of cervical precancer as a predictor of treatment failure: a systematic review and meta-analysis *Lancet Oncol* 2017; 18: 1665–79.

Gatta LA, Kuller JA, Rhee EHJ. Pregnancy outcomes following cervical conization or loop excision procedures. *Obstet Gynecol Surv.* 2017 Aug;72(8):494-499

- Danhof NA, Kamphuis EI, Limpens J, van Lonkhuijzen LR, Pajkrt E, Mol BW. The risk of preterm birth of treated versus untreated cervical intraepithelial neoplasia (CIN): a systematic review and meta-analysis. *Eur J Obstet Gynecol Reprod Biol.* 2015 May;188:24-33.
- D'Alessandro P, Arduino B, Borgo M, Saccone G, Venturella R, Di Cello A, Zullo F. Loop Electrosurgical Excision Procedure versus Cryotherapy in the Treatment of Cervical Intraepithelial Neoplasia: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Gynecol Minim Invasive Ther.* 2018 Oct-Dec;7(4):145-151.
- Silver MI, Gage JC, Schiffman M, Fetterman B, Poitras NE, Lorey T, Cheung LC, Katki HA, Locke A, Kinney WK, Castle PE. Clinical Outcomes after Conservative Management of Cervical Intraepithelial Neoplasia 2 (CIN2) in Women Ages 21-39 Years. *Cancer Prev Res (Phila).* 2018 Mar;11(3):165-170.
- Numnum TM, Kirby TO, Leath III CA, Huh WK, Alvarez RD, Straughn JM. A prospective evaluation of "see and treat" in women with HSIL pap smear results: Is this an appropriate strategy? *J Low Gen Tract Dis* 2005; 9:2-6.
- Li ZG, Qian DY, Cen JM, Chen GD, Shu YH. Three-step versus "see-and-treat" approach in women with high-grade squamous intraepithelial lesion in low-resource country. *Int J Gynecol Obstetrics* 2009;106:202-5.
- Kuroki LM, Bergeron LM, Gao F, Thaker PH, Massad LS. See-and Treat Loop Electrosurgical Excision Procedure for High-Grade Cervical Cytology: Are we Overtreating? *J Low Genit Tract Dis* 2016;20: 247-251.

## **Colposcopy**

- Demarco M, Lorey TS, Fetterman B, Cheung LC, Guido RS, Wentzensen N, Kinney WK, Poitras NE, Befano B, Castle PE, Schiffman M. Risks of CIN 2+, CIN 3+, and Cancer by Cytology and Human Papillomavirus Status: The Foundation of Risk-Based Cervical Screening Guidelines. *J Low Genit Tract Dis.* 2017 Oct;21(4):261-267.
- Stoler MH, Schiffman M; Atypical Squamous Cells of Undetermined Significance-Low-grade Squamous Intraepithelial Lesion Triage Study (ALTS) Group. Interobserver reproducibility of cervical cytologic and histologic interpretations: realistic estimates from the ASCUS-LSIL Triage Study. *JAMA.* 2001 Mar 21;285(11):1500-5.

## **Surveillance**

- Castle PE, Kinney WK, Cheung LC, Gage JC, Fetterman B, Poitras NE, Lorey TS, Wentzensen N, Befano B, Schussler J, Katki HA, Schiffman M. Why does cervical cancer occur in a state-of-the-art screening program? *Gynecol Oncol.* 2017 Sep;146(3):546-553.

Castle PE, Kinney WK, Xue X, Cheung LC, Gage JC, Poitras NE, Lorey TS, Katki HA, Wentzensen N, Schiffman M. Role of Screening History in Clinical Meaning and Optimal Management of Positive Cervical Screening Results. *J Natl Cancer Inst.* 2018 Dec 21

Castle PE, Kinney WK, Xue X, Cheung LC, Gage JC, Zhao FH, Fetterman B, Poitras NE, Lorey TS, Wentzensen N, Katki HA, Schiffman M. Effect of Several Negative Rounds of Human Papillomavirus and Cytology Co-testing on Safety Against Cervical Cancer: An Observational Cohort Study. *Ann Intern Med.* 2018 Jan 2;168(1):20-29. doi: 10.7326/M17-1609.

Gage JC, Hunt WC, Schiffman M, Katki HA, Cheung LA, Myers O, Cuzick J, Wentzensen N, Kinney W, Castle PE, Wheeler CM; New Mexico HPV Pap Registry Steering Committee. Similar Risk Patterns After Cervical Screening in Two Large U.S. Populations: Implications for Clinical Guidelines. *Obstet Gynecol.* 2016 Dec;128(6):1248-1257.

Gage JC, Katki HA, Schiffman M, Castle PE, Fetterman B, Poitras NE, Lorey T, Cheung LC, Behrens C, Sharma A, Zhao FH, Cuzick J, Yang ZH, Kinney WK. The low risk of precancer after a screening result of human papillomavirus-negative/atypical squamous cells of undetermined significance papanicolaou and implications for clinical management. *Cancer Cytopathol.* 2014 Nov;122(11):842-50.

Gage JC, Katki HA, Schiffman M, Fetterman B, Poitras NE, Lorey T, Cheung LC, Castle PE, Kinney WK. Age-stratified 5-year risks of cervical precancer among women with enrollment and newly detected HPV infection. *Int J Cancer.* 2015 Apr 1;136(7):1665-71

Landy R, Cheung LC, Schiffman M, Gage JC, Hyun N, Wentzensen N, Kinney WK, Castle PE, Fetterman B, Poitras NE, Lorey T, Sasieni PD, Katki HA. Challenges in risk estimation using routinely collected clinical data: The example of estimating cervical cancer risks from electronic health-records. *Prev Med.* 2018 Jun;111:429-435.

Schiffman M, Kinney WK, Cheung LC, Gage JC, Fetterman B, Poitras NE, Lorey TS, Wentzensen N, Befano B, Schussler J, Katki HA, Castle PE. Relative Performance of HPV and Cytology Components of Cotesting in Cervical Screening. *J Natl Cancer Inst.* 2018 May 1;110(5):501-508.

### **Risk Modification and Special Populations**

Ahdoot, D., Van Nostrand, K. M., Nguyen, N. J., Tewari, D. S., Kurasaki, T., DiSaia, P. J., & Rose, G. S. (1998). The effect of route of delivery on regression of abnormal cervical cytologic findings in the postpartum period. *Am J Obstet Gynecol*, 178(6), 1116-1120. doi:10.1016/s0002-9378(98)70310-2

Appleby, P., Beral, V., Berrington de Gonzalez, A., Colin, D., Franceschi, S., Goodhill, A., . . . Sweetland, S. (2007). Cervical cancer and hormonal contraceptives: collaborative reanalysis of individual data for 16,573 women with cervical cancer and 35,509 women

- without cervical cancer from 24 epidemiological studies. *Lancet*, 370(9599), 1609-1621. doi:10.1016/s0140-6736(07)61684-5
- Appleby, P., Beral, V., Berrington de Gonzalez, A., Colin, D., Franceschi, S., Goodill, A., . . . Sweetland, S. (2006). Carcinoma of the cervix and tobacco smoking: collaborative reanalysis of individual data on 13,541 women with carcinoma of the cervix and 23,017 women without carcinoma of the cervix from 23 epidemiological studies. *Int J Cancer*, 118(6), 1481-1495. doi:10.1002/ijc.21493
- Bansal, N., Wright, J. D., Cohen, C. J., & Herzog, T. J. (2008). Natural history of established low grade cervical intraepithelial (CIN 1) lesions. *Anticancer Res*, 28(3b), 1763-1766.
- Burchell, A. N., Richardson, H., Mahmud, S. M., Trottier, H., Tellier, P. P., Hanley, J., . . . Franco, E. L. (2006). Modeling the sexual transmissibility of human papillomavirus infection using stochastic computer simulation and empirical data from a cohort study of young women in Montreal, Canada. *Am J Epidemiol*, 163(6), 534-543. doi:10.1093/aje/kwj077
- Burger, E. A., Kim, J. J., Sy, S., & Castle, P. E. (2017). Age of Acquiring Causal Human Papillomavirus (HPV) Infections: Leveraging Simulation Models to Explore the Natural History of HPV-induced Cervical Cancer. *Clin Infect Dis*, 65(6), 893-899. doi:10.1093/cid/cix475
- Castellsague, X., Diaz, M., Vaccarella, S., de Sanjose, S., Munoz, N., Herrero, R., . . . Bosch, F. X. (2011). Intrauterine device use, cervical infection with human papillomavirus, and risk of cervical cancer: a pooled analysis of 26 epidemiological studies. *Lancet Oncol*, 12(11), 1023-1031. doi:10.1016/s1470-2045(11)70223-6
- Cubo-Abert, M., Centeno-Mediavilla, C., Franco-Zabala, P., Merced-Vazquez, C., Castellvi, J., Garcia, A., . . . Xercavins, J. (2012). Risk factors for progression or persistence of squamous intraepithelial lesions diagnosed during pregnancy. *J Low Genit Tract Dis*, 16(1), 34-38. doi:10.1097/LGT.0b013e31822e83cf
- Fader, A. N., Alward, E. K., Niederhauser, A., Chirico, C., Lesnock, J. L., Zwiesler, D. J., . . . Moore, K. N. (2010). Cervical dysplasia in pregnancy: a multi-institutional evaluation. *Am J Obstet Gynecol*, 203(2), 113.e111-116. doi:10.1016/j.ajog.2010.04.016
- Frega, A., Scirpa, P., Corosu, R., Verrico, M., Scarciglia, M. L., Primieri, M. R., . . . Moscarini, M. (2007). Clinical management and follow-up of squamous intraepithelial cervical lesions during pregnancy and postpartum. *Anticancer Res*, 27(4c), 2743-2746.
- Franceschi S, El-Serag HB, Forman D, Newton R, Plummer M. Infectious agents. Thun MJ (ed.). (2018). *Schottenfeld and Fraumeni Cancer Epidemiology and Prevention*. (4th Edition). New York, NY: Oxford University Press. 24: 433-460.
- Fuchs, K., Weitzen, S., Wu, L., Phipps, M. G., & Boardman, L. A. (2007). Management of cervical intraepithelial neoplasia 2 in adolescent and young women. *J Pediatr Adolesc Gynecol*, 20(5), 269-274. doi:10.1016/j.jpog.2007.04.012
- He, Y., Wu, Y. M., Wang, T., Song, F., Wang, Y., & Zhu, L. (2013). Perinatal outcomes of pregnant women with cervical intraepithelial neoplasia. *Arch Gynecol Obstet*, 288(6), 1237-1242. doi:10.1007/s00404-013-2889-6
- Hernandez, B. Y., Wilkens, L. R., Zhu, X., Thompson, P., McDuffie, K., Shvetsov, Y. B., . . . Goodman, M. T. (2008). Transmission of human papillomavirus in heterosexual couples. *Emerg Infect Dis*, 14(6), 888-894. doi:10.3201/eid1406.070616
- Karim, S., Souho, T., Benlemlih, M., & Bennani, B. (2018). Cervical Cancer Induction Enhancement Potential of Chlamydia Trachomatis: A Systematic Review. *Curr Microbiol*, 75(12), 1667-1674. doi:10.1007/s00284-018-1439-7
- Karrberg, C., Brannstrom, M., Strander, B., Ladfors, L., & Radberg, T. (2013). Colposcopically directed cervical biopsy during pregnancy; minor surgical and obstetrical complications and high rates of persistence and regression. *Acta Obstet Gynecol Scand*, 92(6), 692-699. doi:10.1111/aogs.12138

- Koshiol, J., Lindsay, L., Pimenta, J. M., Poole, C., Jenkins, D., & Smith, J. S. (2008). Persistent human papillomavirus infection and cervical neoplasia: a systematic review and meta-analysis. *Am J Epidemiol*, *168*(2), 123-137. doi:10.1093/aje/kwn036
- Mailath-Pokorny, M., Schwameis, R., Grimm, C., Reinthaller, A., & Polterauer, S. (2016). Natural history of cervical intraepithelial neoplasia in pregnancy: postpartum histopathologic outcome and review of the literature. *BMC Pregnancy Childbirth*, *16*, 74. doi:10.1186/s12884-016-0861-8
- Maucort-Boulch, D., Plummer, M., Castle, P. E., Demuth, F., Safaeian, M., Wheeler, C. M., & Schiffman, M. (2010). Predictors of human papillomavirus persistence among women with equivocal or mildly abnormal cytology. *Int J Cancer*, *126*(3), 684-691. doi:10.1002/ijc.24752
- Moscicki, A. B., Hills, N., Shiboski, S., Powell, K., Jay, N., Hanson, E., . . . Palefsky, J. (2001). Risks for incident human papillomavirus infection and low-grade squamous intraepithelial lesion development in young females. *Jama*, *285*(23), 2995-3002. doi:10.1001/jama.285.23.2995
- Onuma, K., Saad, R. S., Kanbour-Shakir, A., Kanbour, A. I., & Dabbs, D. J. (2006). Clinical implications of the diagnosis "atypical squamous cells, cannot exclude high-grade squamous intraepithelial lesion" in pregnant women. *Cancer*, *108*(5), 282-287. doi:10.1002/cncr.22170
- Plummer, M., Peto, J., & Franceschi, S. (2012). Time since first sexual intercourse and the risk of cervical cancer. *Int J Cancer*, *130*(11), 2638-2644. doi:10.1002/ijc.26250
- Rinaldi, S., Plummer, M., Biessy, C., Castellsague, X., Overvad, K., Kruger Kjaer, S., . . . Franceschi, S. (2011). Endogenous sex steroids and risk of cervical carcinoma: results from the EPIC study. *Cancer Epidemiol Biomarkers Prev*, *20*(12), 2532-2540. doi:10.1158/1055-9965.Epi-11-0753
- Rodriguez, A. C., Schiffman, M., Herrero, R., Wacholder, S., Hildesheim, A., Castle, P. E., . . . Burk, R. (2008). Rapid clearance of human papillomavirus and implications for clinical focus on persistent infections. *J Natl Cancer Inst*, *100*(7), 513-517. doi:10.1093/jnci/djn044
- Rositch, A. F., Koshiol, J., Hudgens, M. G., Razzaghi, H., Backes, D. M., Pimenta, J. M., . . . Smith, J. S. (2013). Patterns of persistent genital human papillomavirus infection among women worldwide: a literature review and meta-analysis. *Int J Cancer*, *133*(6), 1271-1285. doi:10.1002/ijc.27828
- Roura, E., Castellsague, X., Pawlita, M., Travier, N., Waterboer, T., Margall, N., . . . Riboli, E. (2014). Smoking as a major risk factor for cervical cancer and pre-cancer: results from the EPIC cohort. *Int J Cancer*, *135*(2), 453-466. doi:10.1002/ijc.28666
- Roura, E., Castellsague, X., Pawlita, M., Travier, N., Waterboer, T., Margall, N., . . . Riboli, E. (2014). Smoking as a major risk factor for cervical cancer and pre-cancer: results from the EPIC cohort. *Int J Cancer*, *135*(2), 453-466. doi:10.1002/ijc.28666
- Schaefer, K., Peters, D., Aulmann, S., Sohn, C., & Eichbaum, M. H. (2012). Value and feasibility of LLETZ procedures for pregnant women with suspected high-grade squamous intraepithelial lesions and microinvasive cervical cancer. *Int J Gynaecol Obstet*, *118*(2), 141-144. doi:10.1016/j.ijgo.2012.02.019
- Schuster, S., Joura, E., & Kohlberger, P. (2018). Natural History of Squamous Intraepithelial Lesions in Pregnancy and Mode of Delivery. *Anticancer Res*, *38*(4), 2439-2442. doi:10.21873/anticancer.12496
- Smith, J. S., Bosetti, C., Munoz, N., Herrero, R., Bosch, F. X., Eluf-Neto, J., . . . Peeling, R. W. (2004). Chlamydia trachomatis and invasive cervical cancer: a pooled analysis of the IARC multicentric case-control study. *Int J Cancer*, *111*(3), 431-439. doi:10.1002/ijc.20257

- Sood, A. K., Sorosky, J. I., Mayr, N., Anderson, B., Buller, R. E., & Niebyl, J. (2000). Cervical cancer diagnosed shortly after pregnancy: prognostic variables and delivery routes. *Obstet Gynecol*, 95(6 Pt 1), 832-838.
- Thomas, D. B., Noonan, L., & Whitehead, A. (1985). Breast cancer and depot-medroxyprogesterone acetate. WHO Collaborative Study of Neoplasia and Steroid Contraceptives. *Bull World Health Organ*, 63(3), 513-519.
- Wetta, L. A., Matthews, K. S., Kemper, M. L., Whitworth, J. M., Fain, E. T., Huh, W. K., . . . Straughn, J. M., Jr. (2009). The management of cervical intraepithelial neoplasia during pregnancy: is colposcopy necessary? *J Low Genit Tract Dis*, 13(3), 182-185. doi:10.1097/LGT.0b013e31818e0c42
- Widdice, L., Ma, Y., Jonte, J., Farhat, S., Breland, D., Shiboski, S., & Moscicki, A. B. (2013). Concordance and transmission of human papillomavirus within heterosexual couples observed over short intervals. *J Infect Dis*, 207(8), 1286-1294. doi:10.1093/infdis/jit018
- Wu, Y. M., Wang, T., He, Y., Song, F., Wang, Y., Zhu, L., . . . Zhang, W. Y. (2014). Clinical management of cervical intraepithelial neoplasia in pregnant and postpartum women. *Arch Gynecol Obstet*, 289(5), 1071-1077. doi:10.1007/s00404-013-3076-5

## New Technologies

- Castle PE, Adcock R, Cuzick J, Wentzensen N, Torrez-Martinez N, Torres S, Stoler M, Ronnett B, Joste NE, Darragh TM, Gravitt PE, Schiffman M, Hunt WC, Kinney WK, Wheeler CM on behalf of the New Mexico HPV Pap Registry Steering Committee<sup>10</sup> and for the p16 Study Panel. Relationships of p16 immunohistochemistry and other biomarkers with diagnoses of cervical abnormalities: implications for LAST terminology *In press* at Arch Pathol Lab Med
- Clarke MA, Fetterman B, Cheung LC, Wentzensen N, Gage JC, Katki HA, Befano B, Demarco M, Schussler J, Kinney WK, Raine-Bennett TR, Lorey TS, Poitras NE, Castle PE, Schiffman M. Epidemiologic Evidence That Excess Body Weight Increases Risk of Cervical Cancer by Decreased Detection of Precancer. *J Clin Oncol*. 2018 Apr 20;36(12):1184-1191. doi: 10.1200/JCO.2017.75.3442. Epub 2018 Jan 22.
- Darragh TM, Colgan TJ, Cox JT, Heller DS, Henry MR, Luff RD, McCalmont T, Nayar R, Palefsky JM, Stoler MH, Wilkinson EJ, Zaino RJ, Wilbur DC; Members of LAST Project Work Groups. The Lower Anogenital Squamous Terminology Standardization Project for HPV-Associated Lesions: background and consensus recommendations from the College of American Pathologists and the American Society for Colposcopy and Cervical Pathology. *J Low Genit Tract Dis*. 2012 Jul;16(3):205-42.